

# SACRIX

*Sacroiliac Fixation*

## *Stop Sacroiliac Joint Pain*



Small incision



Nanofuse used with Sacrix

[www.Nanofusebiologics.com](http://www.Nanofusebiologics.com)



**NANOFUSE  
BIOLOGICS**  
Regenerative Therapeutics



[www.MySacrix.com](http://www.MySacrix.com)

## **Outpatient Spine Surgery**

### **Lateral-Oblique SIJ Transfixing Technique**

### **CPT Code 27279**

Sacrix is inserted ***percutaneously*** using a small incision (~1.5cm) in an ASC with rapid recovery and relief.

REV A

# Sacroiliac Joint Pain

## Symptoms

- Back pain when sitting or laying
- Dull ache below waist
- Buttock pain, may radiate to thigh/groin
- Pain worse climbing stairs or hills

## Potential causes

- Post pregnancy
- Post lumbar fusion
- Sacral fractures
- Arthritis
- Infection
- Instability

## Provocative Tests

### COMPRESSION TEST



### THIGH THRUST TEST



### FABER TEST



### DISTRACTION TEST



### GAENSLER'S TEST



## Surgery Candidates

- Failed medication & therapy for 6 months
- At least 3 positive provocative tests
- X-Ray & CT/MRI imaging
- 75% improvement with image-guided injections
- Steroid injection at least once
- Localized pain over the SI Joint

## Post-Op Instructions

- Keep incision dry for 7-10 days
- Take antibiotics as prescribed
- Take pain meds as prescribed
- Walking as tolerated
- No lifting above 10 lbs
- Light exercises for 8 weeks
- Regular diet
- NO smoking



Nanofuse used with Sacrix

[www.nanofusebiologics.com](http://www.nanofusebiologics.com)

*Sacrix was developed as a collaboration between orthopedics, neurosurgery and interventional pain management.*



Made in USA



**SACRIX**   
Sacroiliac Fixation



*Stop Sacroiliac Joint Pain*